


Medication Dosing Chart

If your child is **less than 2 months old** with a rectal temperature of 100.5 F. or more – call your doctor immediately

The Pediatric Clinic - During office hours: 536-2725 After hours: 395-2200

Acetaminophen (Tylenol)					
Weight:	Milligram Dosage	Infant's Drops 80mg/0.8ml	Children's Liquid/Syrup 160mg/5ml	Children's Soft Chews 80mg each	Junior strength Caps or Chews 160mg each
11-16 lbs	80mg	1 dropper (0.8ml)	½ tsp (2.5ml)	N/A	N/A
17-21 lbs	120mg	1 ½ droppers (1.2ml)	¾ tsp (3.75ml)	N/A	N/A
22-26 lbs	160mg	2 droppers (1.6ml)	1 tsp (5ml)	2 tablets	1 tablet
27-32 lbs	200mg	2 ½ dropper (2ml)	1 ¼ tsp (6.25ml)	2 ½ tablets	
33-37 lbs	240mg	3 droppers (2.4ml)	1 ½ tsp (7.5ml)	3 tablets	1 ½ tablet
38-42 lbs	280mg	3 ½ droppers (2.8ml)	1 ¾ tsp (8.75ml)	3 ½ tablets	
43-53 lbs	320mg	4 droppers (3.2ml)	2 tsp (10ml)	4 tablets	2 tablets
54-64 lbs	400mg	Use Liquid	2 ½ tsp (12.5ml)	5 tablets	2 ½ tablets
65-75 lbs	480mg		3 tsp (15ml)	6 tablets	3 tablets
76-86 lbs	560mg		3 ½ tsp (17.5ml)	7 tablets	3 ½ tablets
87-95 lbs	640mg		4 tsp (20ml)	8 tablets	4 tablets
>95 lbs		Give	Adult	Dosage	

Acetaminophen may be administered every 3 to 4 hours. Ibuprofen (Motrin/Advil) may be administered every 6 to 8 hours
If necessary, acetaminophen and ibuprofen may be alternated.



Medication Dosing Chart

If your child is **less than 2 months old** with a rectal temperature of 100.5 F. or more – call your doctor immediately
 The Pediatric Clinic - During office hours: 536-2725 After hours: 395-2200

Ibuprofen (Advil/Motrin)					
Weight:	Milligram Dosage	Infant Drops 50mg/1.25 ml	Children's Suspension 100mg/5ml	Children's Chewable 50mg each	Junior strength Caps or chews 100 mg each
11-16 lbs		1.25 ml	½ tsp.	NA	NA
17-21 lbs		1.875 ml	¾ tsp.	NA	NA
22-26 lbs		NA	1 tsp.	2 tablets	1 tablet
27-32 lbs		NA	1 1/2 tsp.	3 tablets	1 ½ tablets
33-37 lbs		NA	1 ¾ tsp.	3 ½ tablets	
38-42 lbs		NA	2 tsp.	4 tablets	2 tablets
43-53 lbs		NA	2 ½ tsp.	5 tablets	2 ½ tablets
54-64 lbs		NA	3 tsp.	6 tablets	3 tablets
65-75 lbs		NA	3 ½ tsp.	7 tablets	3 ½ tablet
76-90 lbs			4 tsp.	8 tablets	4 tablets
➤ 90 lbs.		Give	Adult	Dosage	

Acetaminophen may be administered every 3 to 4 hours. Ibuprofen (Motrin/Advil) may be administered every 6 to 8 hours
 If necessary, acetaminophen and ibuprofen may be alternated.